



Review

Facing up to the paradigm of ecological intensification in agronomy: Revisiting methods, concepts and knowledge

Thierry Doré^{a,*}, David Makowski^b, Eric Malézieux^c, Nathalie Munier-Jolain^d, Marc Tchamitchian^e, Pablo Tittone^f

^a AgroParisTech, INRA, UMR 211, F-78850 Thiverval-Grignon, France

^b INRA, AgroParisTech, UMR 211, F-78850 Thiverval-Grignon, France

^c CIRAD, UR HortSys, Boulevard de la Lironde, TA B-103/PS4, 34398 Montpellier Cedex, France

^d INRA, UMR 102 Génétique et Ecophysiologie des Légumineuses, 17 rue Sully, BP 86510, F-21065 Dijon Cedex, France

^e INRA, Unité Ecodéveloppement, Centre PACA, F-84914 Avignon Cedex 9, France

^f CIRAD-Persyst, Systèmes de Culture Annuels, TA B-102/02 Avenue Agropolis, 34898 Montpellier Cedex 5, France

ARTICLE INFO

Article history:

Received 18 October 2010

Received in revised form 5 February 2011

Accepted 16 February 2011

Keywords:

Agroecology
Agroecosystem
Plant science
Farmers' knowledge
Meta-analysis
Comparative analysis

ABSTRACT

Agriculture is facing up to an increasing number of challenges, including the need to ensure various ecosystem services and to resolve apparent conflicts between them. One of the ways forward for agriculture currently being debated is a set of principles grouped together under the umbrella term “ecological intensification”. In published studies, ecological intensification has generally been considered to be based essentially on the use of biological regulation to manage agroecosystems, at field, farm and landscape scales. We propose here five additional avenues that agronomic research could follow to strengthen the ecological intensification of current farming systems. We begin by assuming that progress in plant sciences over the last two decades provides new insight of potential use to agronomists. Potentially useful new developments in plant science include advances in the fields of energy conversion by plants, nitrogen use efficiency and defence mechanisms against pests. We then suggest that natural ecosystems may also provide sources of inspiration for cropping system design, in terms of their structure and function on the one hand, and farmers' knowledge on the other. Natural ecosystems display a number of interesting properties that could be incorporated into agroecosystems. We discuss the value and limitations of attempting to ‘mimic’ their structure and function, while considering the differences in objectives and constraints between these two types of system. Farmers develop extensive knowledge of the systems they manage. We discuss ways in which this knowledge could be combined with, or fed into scientific knowledge and innovation, and the extent to which this is likely to be possible. The two remaining avenues concern methods. We suggest that agronomists make more use of meta-analysis and comparative system studies, these two types of methods being commonly used in other disciplines but barely used in agronomy. Meta-analysis would make it possible to quantify variations of cropping system performances in interaction with soil and climate conditions more accurately across environments and socio-economic contexts. Comparative analysis would help to identify the structural characteristics of cropping and farming systems underlying properties of interest. Such analysis can be performed with sets of performance indicators and methods borrowed from ecology for analyses of the structure and organisation of these systems. These five approaches should make it possible to deepen our knowledge of agroecosystems for action.

© 2011 Elsevier B.V. All rights reserved.

Contents

1. Introduction.....	198
2. Diversifying sources of knowledge to guide ecological intensification.....	199
2.1. Mobilizing advances in plant sciences.....	199

* Corresponding author. Tel.: +33 1 30 81 52 45; fax: +33 1 30 81 54 25.

E-mail address: thierry.dore@agroparistech.fr (T. Doré).

2.1.1.	A new look at the basics	199
2.1.2.	The cultivated plant and its biological environment	199
2.1.3.	Ways to improve the use of plant sciences for ecological intensification	200
2.2.	Learning lessons from the functioning of natural ecosystems	201
2.2.1.	What does “Mimicking natural ecosystems” mean?	201
2.2.2.	Agroecosystems as complex socio-ecological systems	202
2.3.	Farmers’ knowledge and lay expertise valorisation and integration into scientific knowledge	203
2.3.1.	Value of farmers’ knowledge for agronomy	203
2.3.2.	Qualification and validation of lay expertise and knowledge expression	204
3.	Methods for synthesizing information	204
3.1.	Meta-analysis and agronomy	205
3.2.	Comparative analysis of agroecosystems	205
3.2.1.	Comparative analysis based on multiple indicators	206
3.2.2.	Analysing the structure and functioning of agroecosystems	206
4.	Overall discussion and conclusion	207
	Acknowledgement	208
	References	208

1. Introduction

New agricultural systems are required to allow agriculture to satisfy the increasingly diverse expectations of society. For decades, agronomy has produced knowledge and designed agroecosystems for maximising the production of primary food and fibre, either for direct consumption or for industrial use. Agricultural production issues have recently been expanded to include other ecosystem services (Zhang et al., 2007). Like other natural and semi-artificial ecosystems, agroecosystems can provide services, such as carbon sequestration, pollination, or water filtration. The capacity of agriculture to provide such services is, of course, not always guaranteed, and there are many examples of adverse effects of agricultural practices on the environment, leading to ecological disservices of agriculture (Matson et al., 1997; Swinton et al., 2007). Disservices may include decreases in water and air quality or a contribution to biodiversity loss. As agroecosystems are ecosystems controlled by humans, adopting the correct approach to a wide range of production issues requires an understanding of the way in which natural and human-driven or forced processes interact within the ecosystem.

Agronomists have argued that the missions of multi-objective agriculture could best be achieved by making better use of biological regulation mechanisms at different levels: crop management, cropping system design, landscape layout and management (Matson et al., 1997; Médiène et al., 2011). This assumes that biological mechanisms are able to replace chemical or physical inputs, or to interact favourably with them, playing the same agronomic role without external costs, including environmental costs in particular. The use of biological regulation in agroecosystems to achieve both a high level of food production and to provide ecosystem services, apparently opposite aims, has been placed at the core of what is increasingly called “ecological intensification”. The Food and Agriculture Organisation (FAO, 2009) recently defined “ecological intensification” (or “sustainable intensification”) within the framework of organic agriculture as “Maximization of primary production per unit area without compromising the ability of the system to sustain its productive capacity”. The expression “ecological intensification” was already in use more than two decades ago (Egger, 1986), when it referred to a kind of ecological engineering in agropastoral systems in Africa, replacing some perennial species to improve soil organic matter content.

A more recent use of the expression by Cassman (1999) focused on cereal production and highlighted the need for progress in plant and soil science to achieve a continuous increase in cereal yields (intensification) without environmental (ecological) dam-

age. This approach focuses principally on the fate of fertilisers and their use by crops. Witt et al. (2006) applied a similar approach to oil palm plantations. According to Chevassus au Louis and Griffon (2008) and a number of other authors (Affholder et al., 2008; Mikolasek et al., 2009; Hubert et al., 2010; Bommel et al., 2010), ecological intensification is a pathway towards the production of more agricultural product, the production of “new” things (ecosystem services) and different means of production (environmentally friendly). According to Chevassus au Louis and Griffon (2008), ecological intensification is based on “intensification in the use of the natural functionalities that ecosystems offer”. Though relatively vague, this definition remains a possible starting point for the consideration of alternative pathways of development for agriculture. This definition is much broader than that of Cassman (Cassman, 1999), and provides an interesting haven for scientists promoting the use of biological regulation in agroecosystems.

Many articles have been published on biological regulation in agroecosystems, mostly under the heading “agroecology”, and new papers are continuing to appear. Research on this topic remains highly necessary, and is probably a challenge for most agronomists familiar with individual physical and/or chemical aspects of agroecosystems. However, ecological intensification calls for both a wider diversification of sources of knowledge and the development of new data analysis methods. Agronomists have, until recently, relied essentially on their own scientific output. Prototyping (e.g., Vereijken, 1997; Lançon et al., 2007; Debaeke et al., 2009) and the model-based design of agricultural systems (e.g., Rossing et al., 1997; Bergez et al., 2010) are fed by results processed through simulation studies, statistical hypothesis testing and group analysis, from research groups working mostly at experimental stations (Fig. 1). We argue here that agronomists would be placed in a better position to tackle ecological intensification if they diversified their sources of knowledge and the methods used to compile, organise and analyse such knowledge. The diversification of knowledge sources may include (i) making use of recent advances in plant sciences, (ii) learning lessons from the functioning of natural ecosystems, guiding the design and management of agroecosystems and (iii) embracing local farmers’ knowledge. Methods for assessing these sources of knowledge are necessarily diverse, and could be extended to data mining and the meta-analysis of large datasets containing heterogeneous information and comparative analyses of agroecosystems at different scales. We present here the arguments for further agronomic research in these two related domains: sources of knowledge for agronomists and data processing methods.




Fig. 1. Summary of new avenues of agronomic research for ecological intensification.

2. Diversifying sources of knowledge to guide ecological intensification

2.1. Mobilizing advances in plant sciences

There has been tremendous progress in plant sciences in recent decades, with detailed elucidation of the genetic and environmental determinism of plant development, growth and reproduction. This progress was made possible, in particular, by increases in our ability to dissect cellular and molecular processes, supported by exponential progress in laboratory techniques and the capacity to analyse masses of genomic data (e.g., Tardieu and Tuberosa, 2010). This knowledge about the highly complex life of plants has often been developed in a simplified environment, far removed from the reality of farmers' fields. This has led to a widening of the gap between the research objectives of plant scientists and agronomists. We highlight briefly, with a few examples, ways in which agronomists could make use of advances in plant sciences to design ecologically intensive cropping systems.

2.1.1. A new look at the basics

Agronomists involved in the design and evaluation of cropping systems often make use of a simplified crop description (Monteith, 1977), despite the availability of more mechanistic models simulating canopy photosynthesis (Spitters et al., 1986; Spitters, 1986; dePury and Farquhar, 1997). In this simplified description, the canopy, represented as a "big leaf", intercepts photosynthetically active radiation and converts it to biomass. Branching is generally considered to be the outcome of interplant competition. Mineral nutrition is represented as a simple flux from soil to plant roots, depending on soil mineral and water contents. Such simplified representations have proved sufficient and highly successful for cropping system design. Moreover, the more sophisticated representations of the basic processes of plant life implemented in more complex models do not necessarily improve the ability of crop models to predict behaviour in a range of fluctuating conditions. Such representations have therefore been used only rarely

by agronomists. Nevertheless, results recently obtained in plant sciences suggest that this simple paradigm could be improved, as shown for example by Zhu et al. (2010), who analysed the ways in which improvements in photosynthesis efficiency could contribute to the required increase in yields.

Nutrient use efficiency is also clearly a key point in ecological intensification. One of the most important issues is decreasing the use of nitrogen fertilisers, to decrease greenhouse gas emissions, to reduce the dependence of agriculture on fossil fuels and to prevent health and environmental disorders, without decreasing productivity (Galloway et al., 2008; Spiertz, 2010). Plant scientists have investigated in detail the exchanges of nitrogen between roots and their environment (Jackson et al., 2008). Glass (2003) summarised the factors decreasing nitrogen absorption efficiency, on the basis of molecular knowledge and empirical data. Decreases in nitrogen transporter activity and rates of nitrate absorption follow increases in soil ammonium concentration, low temperature and incident radiation. These mechanisms may account, at least in part, for the high variability of fertiliser efficiency observed in field experiments. They also provide us with opportunities to improve nitrogen management in the soil. More generally, the ways in which plants make use of adaptation mechanisms to deal with mineral depletion have been extensively studied on a physiological basis (Grossman and Takahashi, 2001). Agronomists could make use of this work to define the limits within which plant environments must be contained to avoid unfavourable plant reactions.

2.1.2. The cultivated plant and its biological environment

Since the middle of the last century, the gradual "artificialisation" of agriculture has led to agronomists paying less attention to the biological components of fields. Agroecology has emerged as a reaction against this excessive simplification of the system, placing the biological component back at the heart of the system (Altieri, 1989), and resulting in the development of an "agroecosystem" view (Conway, 1987). Nevertheless, common agronomic practices still largely ignore biological interactions in cultivated fields, and agroecologists often emphasise the need for an empirical and

holistic approach to agroecosystems. New findings in plant sciences concerning the relationships between the plant and its surrounding biotic environment have recently emerged and are of great interest.

Studies of interactions between roots and soil micro- and macro-organisms have revealed the existence of processes of paramount importance for agronomists. Some of these interactions are very familiar to agronomists, including nitrogen fixation by symbiosis between *Rhizobium* sp. and leguminous or non-leguminous (Mehboob et al., 2009) plants. Other associations, such as that between other endophytic di-azotrophic bacteria and grasses or cereals, also exist and may be of interest, as pointed out by Reis et al. (2000). Plants may be injured by soil pathogenic organisms, but they may also benefit from organisms present in the rhizosphere, through improvements in growth and mineral nutrition, an increase in resistance to unfavourable abiotic conditions, and protection against or an increase in resistance to pathogens (Sturz and Nowak, 2000; Kiers and Denison, 2008).

Whatever the types of organisms considered, the species or plant genotype drives selection of the bacterial community and determines the benefits of plant–rhizosphere mutualism. Improvements in the genomic characterisation of rhizobacterial communities have made it possible to demonstrate that plant genotype influences bacterial assemblages by modifying exudation patterns (Micallef et al., 2009). An understanding of the plant genome would make it possible to determine the genetic basis of the mechanism and to make use of genetic variants for the management and manipulation of the rhizosphere community (Ryan et al., 2009; Wissuwa et al., 2009). These rhizosphere associations and their benefits to the crop also depend strongly on cropping system, so it would seem reasonable to conclude that adapted cropping systems (including crop rotation and crop management measures) could also increase efficiency. The efficacy of the *Rhizobium*/legume association is also highly dependent on cropping system, through the effects of practices on the physical and chemical properties of soils and their water status (Sprent et al., 1987). These effects are well known, but should be considered in the light of the recent development of legume nodulation genomics (Stacey et al., 2006). Sturz and Nowak (2000) have enlarged their vision to the overall communities of endophytic rhizobacteria with potentially beneficial effects on crop growth through an increase in resistance to unfavourable abiotic conditions and to pathogen aggression, and through improvements in growth and mineral nutrition. The agronomic benefits of these associations with endophytic rhizobacteria depend on the survival of bacterial communities, which in turn depends on soil and crop management (Bowen and Rovira, 1999; Acosta-Martinez et al., 2008). One of the ways by which crop management can modulate the evolution of microbial communities, is its effect on root exudates. In addition to altering the physical and chemical properties of the soil, root exudates have been shown to affect both soil micro-organism communities and other eukaryotes (Bertin et al., 2003). Bais et al. (2004, 2006) reviewed the nature of the chemicals involved and the corresponding interaction processes for various ecological roles. However, one of the aspects of crop/soil community interactions most frequently ignored by agronomists is probably the role of the common mycorrhizal networks (CMNs), which may be affected directly or indirectly by soil tillage, fertilisers, pesticide use and aerial plant management (Pietikäinen and Kytöviita, 2007). The networks that these fungi establish between plants may provide a major route for mineral transfer from plant to plant (He et al., 2003). van der Heijden and Horton (2009) recently reviewed the possibilities for CMN formation between different plant species, their ecological significance and the benefits generated. They found that there were many possibilities for CMN development, but that there were also large differences in the benefits accrued, particularly in terms of promotion of the growth of interconnected plants. Similarly, the role

of plant micro-organisms in plant × plant interactions (Sanon et al., 2009; Li et al., 2008) and the competition of microbial communities promoting both plant growth and health (Lemanceau et al., 2009) illustrate the benefits that agronomists may obtain from advances in research on plant–micro-organism interactions for rhizosphere engineering and management (Ryan et al., 2009). Beyond the question of production, Jackson et al. (2008), focusing on nitrogen, derived from current knowledge on root/micro-organism interactions the trends in ecosystem services supplied by cropping systems in different agricultural situations. Thanks to the deep insight now available, the contribution of agronomists at system level can be built on mechanistic rather than empirical knowledge, as demonstrated by certain examples in precision agriculture (Welbaum et al., 2004).

Interactions between aerial parts of the plant and the surrounding biotic environment have also been described in detail in recent years. The metabolic pathways by which plants react both locally and systemically to infection or wounding are increasingly well known (de Bruxelles and Roberts, 2001; Kessler and Baldwin, 2002). Some result in the production of volatile substances, which play a role in herbivore repulsion or plant-to-plant signalling. These findings are promising for genetic engineering approaches, provided that the genetic basis of the metabolic pathways can be identified (Dudareva and Pichersky, 2008). However, cropping system may also play a role, as the expression of the metabolic pathways involved in direct or indirect defence probably depends on interactions between genotype and environment (Le Bot et al., 2009). Moreover, it may be possible to elicit some of these pathways deliberately, with appropriate techniques.

2.1.3. *Ways to improve the use of plant sciences for ecological intensification*

The preceding two sections do not provide a detailed review of the extensive literature in plant sciences. Instead, they deal with a few examples of recent progress and the possible benefits that agronomists could derive from these advances (see Table 1). These examples demonstrate that closer consideration of the results of plant sciences could help agronomists to reach their objectives, paving the way for higher levels of production, better quality products, and less harmful consequences for the environment. Other advances in plant sciences, concerning plant architecture, leaf and root morphogenesis (McSteen and Leyser, 2005; Wang and Li, 2008; Walter et al., 2009), floral biology (e.g., Boss et al., 2004), the role of aquaporins (e.g., Maurel et al., 2008), cell separation processes (Roberts et al., 2002) and long distance signals within plants (Lough and Lucas, 2006), for example, are also of great potential interest to agronomists working on ecological intensification, as they might help crops to avoid or to resist deleterious stresses. However, major efforts are still required to scale-up the results from individual genes, cells or organs to the canopy, and to test the stability of biological results in a wide range of agricultural conditions. It is also important to check that advances in one area are not associated with severe drawbacks in others. However, these findings are nonetheless precious to agronomists, who will need to use all the means available to construct novel, more resource-use efficient and/or productive cropping systems.

Finally, there are many different drivers of change in ecological intensification (see introduction and subsequent sections). Innovative systems that have already been developed in the domain of ecological intensification, such as the use of mixtures of cultivars or species, agroforestry and no-tillage systems, would certainly benefit from the knowledge provided by plant sciences. However, these systems will themselves raise new questions and issue new challenges to plant science. For example, although progress has been made in this area, plant sciences

Table 1
Examples of recent results from plant sciences useful in agronomy.

Topics in plant sciences	Key references	Potential agronomic benefits
Plant architecture	Zhu et al. (2010) Walter et al. (2009) dePury and Farquhar (1997)	Increased radiation interception
Photosynthesis efficiency	Wang and Li (2008)	Canopy pattern target for crop management Increase in yield Identification of genotypes adapted for crop mixture
Exchanges of nitrogen between roots and environment Role of organic anion exudation	Jackson et al., 2008 Glass (2003) Ryan et al. (2001)	Improved fertiliser use efficiency Improved nitrogen management
Interaction between roots and soil organisms	Mehboob et al. (2009) Brussaard et al. (2007)	Improved mineral nutrition
Role of common mycorrhizal networks	Micallef et al. (2009) Ryan et al. (2009) Sturz and Nowak (2000) van der Heijden and Horton (2009)	Improved crop growth Adaptation of crop management
Interaction between aerial parts of the plant and environment	de Bruxelles and Roberts (2001)	Management of natural defences for improved resistance to pests

results are still often obtained in highly simplified systems and therefore cannot easily be translated to multispecies systems. Above-ground competition for light and below-ground competition for water are major processes in ecological intensification that require study in systems including facilitation between plants (Long and Nair, 1999; Zhang et al., 2008; Malézieux et al., 2009).

2.2. Learning lessons from the functioning of natural ecosystems

Strategies for agroecosystem design and management may be derived from the observation of natural ecosystems, guiding alternative agronomic practices. Several authors (e.g., Ewel, 1999; Altieri, 2002; Jackson, 2002; Vandemeer, 2003; Malézieux, 2011) have already suggested that natural ecosystems may provide appropriate models for agroecosystem design to achieve both environmental and social goals while ensuring long-term sustainability. This idea is based on the assumption that natural ecosystems are adapted to local constraints, due to a long process of natural selection (Dawson and Fry, 1998; Ewel, 1999). It is therefore assumed that the incorporation of certain characteristics of natural ecosystems into agroecosystems would improve some of the properties of agroecosystems, such as productivity (Fukai, 1993), stability (Aerts, 1999; Schulte et al., 2003) and resilience (Lefroy et al., 1999). These features are particularly useful for dealing with pest outbreaks (Trenbath, 1993) and increasing energy efficiency in a context of the depletion of fossil fuels (Hatfield, 1997). A similar reasoning was followed in the framework of Ecoagriculture, proposed by McNeely and Scherr (2003), which places biodiversity at the heart of strategies to conserve and restore ecosystem services, increase wild populations in agroecosystems, and sustain agricultural production. An illustration of this mimicry is provided for cropping systems in Fig. 2 with an emphasis on crop protection. In natural ecosystems, the various animal and plant species interact through population dynamics and trophic networks, providing the final ecosystem with services, such as pollination. In standard cropping systems, these interactions may lead to pest damage on crops, which may be managed with various control methods to limit yield loss. An increase in plant species diversity in systems mimicking natural ecosystems could allow natural enemies to control pests and generate ecosystem services.

2.2.1. What does “Mimicking natural ecosystems” mean?

There have been only a few practical attempts to design agroecosystems from nature. Jackson and Jackson (1999) aimed to develop sustainable cropping systems by mimicking the mid-grass American prairie, creating crop mixtures analogous to the vegetation structure of the prairie. Traditional agroecosystems in the tropics, long unknown or disparaged by some agronomists, are frequently based on the integrated management of local natural resources and, in many cases, on the management of local biodiversity. These systems may also be considered to result from the observation of nearby natural ecosystems by generations of farmers, who have aimed to mimic the functioning and structure of these natural systems. For example, slash and burn systems can be considered to mimic nature behaviour after fire. Agroforestry systems in the humid tropics mimic the structure and functioning of rainforests. According to Ewel (1999), humid tropical ecosystems appear to be particularly suitable for application of the “mimicry of Nature” concept. Agroforestry systems in the humid tropics are based on the tropical rainforest model. They combine several strata, have a high level of species diversity and are very widespread in Asia, Oceania, Africa and Latin America. Such systems provide both subsistence for local populations and major environmental and socio-economic services (Sanchez, 1995; Nair, 2001). Lying halfway between agro- and forest ecosystems, agroforestry systems combine annual and perennial, herbaceous and woody species, in a more or less complex whole in terms of the number of plant species and practices (Torquebiau, 2007). The damar agroforests of Sumatra, or the cocoa-based agroforests of Cameroon or Costa Rica, are original ways in which farming communities use natural resources in human reconstructions of both “natural” and productive ecosystems from natural ecosystems (Michon et al., 1995, 2007; Schroth et al., 2001, 2004).

The scientific foundations of the mimicry paradigm, however, remain to be studied thoroughly (Malézieux, 2011). The potential of this approach to generate innovative agroecosystems in practice also remains largely unknown. Ewel (1999) and van Noordwijk and Ong (1999) proposed two principles for the design of agroecosystems based on natural ecosystem mimicry. According to the first of these principles, agroecosystems should mimic the structure and function of natural ecosystems existing in a given pedoclimatic zone. According to the second, agroecosystems should also mimic




Fig. 2. A comparison of natural ecosystems, conventional cropping systems and agroecosystems inspired from natural ecosystems, with an emphasis on crop protection.

the diversity of species existing in natural ecosystems, thereby maintaining the diversity of natural ecosystems in the given zone. The first of these principles is clear enough, but must be extended to be effective. Indeed, there are many functions, and structure can be assessed at different scales. Furthermore, basing agroecosystem design solely on natural ecosystems present in the same area may be too limiting: some good ideas might emerge from the study of very distant systems.

According to the second principle, the redesign of agroecosystems in more ecologically intensive configurations implies their diversification. This has been the case, for example, in Cuba, where small- and medium-scale farmers have tended to diversify their production systems in response to their limited access to or total lack of agricultural inputs to sustain productivity (Funes-Monzote et al., 2009). The resulting diversified systems are energetically more efficient, less dependent on external inputs, more productive, adaptable and resilient. The diversification of agroecosystems within the mimicry paradigm may be achieved by increasing the number of microorganisms, plant and animal species relevant to agriculture over space and time, or through agrobiodiversity, a subset of general biodiversity (Brookfield et al., 2003). However, natural ecosystem mimicry cannot mean reproducing the diversity observed in natural ecosystems, for at least three reasons. First, recent reviews of existing knowledge in ecology have demonstrated that functional composition controls ecosystem functioning more frequently than species diversity (Hooper et al., 2005). As our purpose is to improve agroecosystem functioning through ecological intensification, and not to conserve natural species biodiversity *per se* within agroecosystems, agronomists should concentrate on identification of the level of functional biodiversity resulting in the expression of interesting properties. As pointed out by Main (1999), who addressed the question of how much biodiversity is enough in the context of agroecosystems mimicking nature, the level of diversity considered adequate strongly depends on the goals and criteria used for evaluation. Moreover, interesting properties may arise from the spatial and temporal organisation of the species rather than purely from their number. For example, lessons can be learned from studies of natural ecosystems addressing agronomic topics: nutrient cycling within a complex landscape may be useful

for optimising nutrient management in areas worked by humans, community ecology in natural ecosystems may facilitate the design of new crop protection strategies and an understanding of facilitation within natural ecosystems should make it easier to make use of this process in agroecosystems. Finally, approaches based on mimicking natural ecosystems will inevitably be confronted with the "aim problem". Natural ecosystems provide many services but are not targeted. Agroecosystems, by contrast, are designed to optimise different aspects and to achieve different goals. Consequently approaches mimicking natural ecosystems are limited by certain agricultural obligations, such as the removal of the minerals contained in agricultural products. Some insight may be gained from regarding agroecosystems as complex systems with many simultaneous feedback loops including a dimension absent from natural ecosystems: human agency.

2.2.2. Agroecosystems as complex socio-ecological systems

Agroecosystems are systems that combine sociological and ecological dynamics, in interaction. In complex, dynamic and spatially heterogeneous systems, interactions take place over scales generating emergent properties and self-regulatory mechanisms (Holling, 1973). These mechanisms often manifest as cross-scale feedback, or *panarchy* (Gunderson and Holling, 2002), and societies contribute to system regulation through adaptive management. For example, in smallholder agricultural systems making use of communally shared resources, buffering and regulatory mechanisms often emerge from collective action (Meinzen-Dick et al., 2004). This is why agroecosystems may be defined as socio-ecological systems, or cybernetic systems steered by humans to attain certain goals (see Conway, 1987). The capacity of farmers to adapt plays a major role in system resilience and, by analogy to the concept of informal economies (de Soto, 2000), regulatory mechanisms operate as informal resource flows that are often unaccounted for in agroecosystems analysis (Tittonell et al., 2009). Just as natural ecosystems have a "memory" as a direct consequence of their history, so do agroecosystems, except that some of that memory lies in human agency (Tittonell, 2007).

A wider definition of agroecosystem diversification, more compatible with the socio-ecological nature of complex

agroecosystems, must consider not only species diversity, but also the diversity of agricultural practices and rural knowledge adapted to/derived from local pedoclimatic conditions. These lie at the core of human agency and represent new sources of knowledge for agronomic research (see below). Agroecosystem diversification in its broadest sense thus concerns the diversity of livelihood strategies at a certain location, diverse land use, management and marketing strategies, the integration of production activities (e.g., crop-livestock interactions), spatial and temporal associations of crops and crop cultivars, and the maintenance of genetic agrobiodiversity in the system. The efficiency of use of natural, economic and social resources in agroecosystems—which goes beyond the partial use efficiency of a certain single input—and desirable properties, such as stability and resilience, are based on one or more of these categories of diversity. New avenues for agronomy to strengthen agroecological intensification should go beyond the cultivated field or the mixture of species in a given landscape. They should explore desirable properties and mechanisms that operate at the scale of complex socio-ecological systems, i.e. that take into account sociological and ecological dynamics and interactions in agroecosystems.

2.3. Farmers' knowledge and lay expertise valorisation and integration into scientific knowledge

Farmers do not rely exclusively on the results and output of agronomic research to operate their agroecosystems. They make use of much wider knowledge, based on their own experiences and on exchanges with other farmers and advisers, thus building their own expertise. This expertise is rooted in the need to act whatever the level of agronomic knowledge available: sound and detailed or unreliable and patchy. It is also dependent on the characteristics (environmental, economic, social) of the situation in which it is constructed. According to [Prior \(2003\)](#), we may consider farmers to be *lay experts* (although this denomination entails an antinomy): *experts* because of their experience-based knowledge and *lay* because this knowledge is limited in scope and does not give farmers the broader and deductive understanding characteristic of scientific or expert knowledge. Recognition of the value of lay expertise is both a necessity and a challenge in many domains, such as medicine (e.g., adapting treatments according to the patient's reactions, both as observed by doctors and as interpreted by the patient) and industry (particularly for fault detection in plant or machine operation). However, although the value of this lay expertise is recognised, it is not used to build or extend the current scientific knowledge, but to adapt its application in local situations ([Henderson, 2010](#)).

Farmers can observe not only their own production systems, but also other systems (both agricultural and natural) and interactions between these systems. They can also gain experimental knowledge in their own systems. They are often willing to do so and therefore carry out experiments in the operation of their own agroecosystem, evaluating the response of the system to their decisions. This generates different types of knowledge. When confronted with, observing or learning from natural ecosystems, farmers gain knowledge similar to what is generally referred to as *local or traditional ecological knowledge* (LEK or TEK, [Berkes, 1999](#)). Over generations, they may also build traditional knowledge (not specifically ecological), refined by years of adaptation (see previous section). When experimenting, they build a mixture of experience-based and experimental knowledge. Many studies have considered the use of LEK/TEK, but most have focused on the use of this knowledge for natural resource management (including fisheries and forestry systems, which more closely resemble a subsistence harvesting activity) rather than the design or improvement of productive agricultural systems. Fewer studies have directly

Table 2
Examples of farmers' knowledge potentially useful in agronomy.

Sources of knowledge	Key references	Potential agronomic benefit
Local ecological knowledge	Chalmers and Fabricius (2007)	Explaining changes in agricultural systems
Traditional farming systems	Singh and Sureja (2008) Abbona et al. (2007)	Design of sustainable farming systems Understanding of ecological processes
Local knowledge and indicators for assessing forest management	Ballard et al. (2008)	Assessment of management practices for forests
Farmer's indicators supporting decision making	Tchamitchian et al. (2006)	Indicators with expanded domains of validity

investigated farmers' knowledge. The studies that have been carried out in this domain have mostly assessed the validity of this knowledge (e.g., [Grossman, 2003](#); [Friedman et al., 2007](#); [Grace et al., 2009](#)) or considered the local adaptation of more generic solutions (e.g., [Steiner, 1998](#); [Affholder et al., 2010](#)). However, farmers' knowledge is not only of value for application and for the adaptation of agronomic knowledge to a particular case. It can also be used to extend the available scientific agronomic knowledge (see the examples presented in [Table 2](#)). We will defend this point and discuss the various issues it raises below.

2.3.1. Value of farmers' knowledge for agronomy

We will analyse separately the lay expertise (resulting from farmers' activities and interactions with their own systems) and the more traditional knowledge that some farmers or societies have developed over time. The value of lay expertise for agronomy and for development (support to farmers) has been recognised for some time (e.g., [Barzman et al., 1996](#); [Baars and de Vries, 1999](#)). This lay expertise can help to enlarge current agronomic knowledge in various ways. First, farmers operate their agroecosystem even in the absence of appropriate knowledge, because they have to. They therefore develop experience-based knowledge that can fill in some of the gaps in scientific knowledge. However, as mentioned above, this experience-based knowledge is often limited to the farmer's own particular case, whereas scientific knowledge should be more general.

Second, some traditional practices are based on the observation of natural ecosystems ([Chalmers and Fabricius, 2007](#); [Reed et al., 2007](#)), which, as we have seen, may be of value for ecological intensification. [Chalmers and Fabricius \(2007\)](#), for example, showed that local experts, using their ecological knowledge, were able to put forward explanations for changes in their system, some of which were also provided by scientific knowledge. However, the local experts also had other explanations rooted in a more general understanding of the system. Traditional farming systems can also be a source of understanding and inspiration for the design of sustainable farming systems. [Singh and Sureja \(2008\)](#) showed, for example, how traditional farming systems cope with harsh environments through the management of a wide diversity of plants providing genetic resources. [Abbona et al. \(2007\)](#) evaluated the sustainability of a traditional vineyard system in Argentina, both in its original location and in a newly planted area. They showed that the traditional system, in its original location, was indeed sustainable, whereas this system was not sustainable in its new, different location. They concluded that the efficacy of the traditional system was dependent on the location in which and for which it had been developed over time. During this evaluation process, based on the use of indicators developed for this analysis through

the adaptation of existing methods, these authors gained insight into and an understanding of the ecological processes at work in the traditional vineyard system. The analysis of traditional farmers' practices therefore provided an opportunity to obtain new scientific knowledge. In a different context, Ballard et al. (2008) analysed the knowledge involved in the management and monitoring activities of community-based forestry groups and the ways in which local and scientific knowledge complemented each other. They showed that local knowledge provided a rapid and efficient means of assessing the effects of management practices on the forest. The same was found for greenhouse tomato management. Tchamitchian et al. (2006) successfully used the concept of "crop vigour" as an indicator in their expert system controlling the daily greenhouse climate for tomato production. Tomato crop vigour is readily assessed by growers of greenhouse tomato crops, on the basis of a set of observations: plant tip colour and shape, fruit load on the crop, crop overall colour. Scientists relate these observations to the generative to vegetative balance of the crop and its ability to perform photosynthesis (Navarrete et al., 1997), without being able to model it formally.

Taken as a whole, local knowledge and lay expertise can provide clues to the natural or ecological processes most useful in the design of sustainable farming systems, such as the natural regulation of pest populations by their predators (Barzman et al., 1996; Sinzogan et al., 2004), or management of the soil and its mineral balance (Steiner, 1998; Okoba and de Graaff, 2005; Saito et al., 2006; Abbona et al., 2007). They can also be of value in the design of assessment methods or indicators for monitoring the ecological performances of these farming systems.

2.3.2. Qualification and validation of lay expertise and knowledge expression

Although both interesting and challenging, the lay expertise of farmers (or advisers) is not easy to use. First, this lay expertise must be elicited and represented. Several methodologies have been proposed for expert knowledge elicitation, either for specific applications, such as plant disease epidemics (Hughes and Madden, 2002), or for more general applications (Cornelissen et al., 2003; Ley et al., 2010). Appropriate elicitation methods include the selection of a panel of experts and the associated delimitation of the knowledge domain considered. The choice of representation also influences the elicitation process. Many authors advocate the use of fuzzy models, which allow the use of linguistic terms and are more suitable for the expression of knowledge in qualitative rather than quantitative terms. By contrast, scientific knowledge is most frequently modelled in quantitative terms, particularly when the goal is to represent the operation of a system under the influence of both controlled (human decisions and actions) and uncontrolled (environment) factors. Most of the agronomic models built to simulate agroecosystems are numerical models in which the variables have point values rather than interval or probabilistic values. There is therefore a gap between the most common representation of scientific knowledge and that of lay expertise, hindering the combination and merging of these two types of knowledge. However, differences in representation are not the only difficulty. As pointed out by Prior (2003), lay experts may be wrong, either because of the limited scope of their experience or because their conclusions are based on false premises (misobservations, for example, due to a lack of knowledge or skills). Their knowledge is also situation-dependent in that it is obtained in a domain of low variability (one of the goals of agricultural practices is often to reduce variability and diversity in agroecosystems, a goal challenged by ecological intensification). Lay expertise should therefore be qualified and analysed independently, in several different ways: domain of validity, certainty and precision. The domain of validity is important because knowledge should be associated with a description of the domain

in which it was obtained (ranges of the variables considered, for example); this factor can be used to analyse the extent to which the knowledge obtained is generic. Certainty refers to the confidence that can be attributed to the knowledge. Finally, precision measures how close to a numerical expression it is possible to get in the expression of the knowledge. Even certain knowledge may display a low precision rendering its use purely hypothetical (ventilating a greenhouse does modify its temperature, but the change is difficult to indicate with precision). Artificial intelligence provides a framework for representing expertise and analysing the conflicts arising when information from different sources is compared (several lay experts or a combination of lay expertise and scientific knowledge; Amgoud and Kaci, 2007; Bench-Capon and Dunne, 2007; Alsinet et al., 2008; Amgoud and Prade, 2009). However, this domain (qualitative reasoning and argumentation) is still developing and, to our knowledge, its concepts and tools have not yet been used to merge lay expertise and scientific knowledge in agronomy (there are applications for database fusion, assisting debate preparation and industrial planning). The added value of these approaches lies in the need to provide an explanation detailing the arguments supporting a piece of knowledge, therefore addressing the questions of certainty and precision raised above.

The qualification of lay expertise has been shown to be a necessary step in approaches aiming to combine this expertise with scientific knowledge. Going beyond the issues of the domain of validity, certainty and precision, there is the question of validation of the new knowledge obtained. However, classical validation procedures cannot readily be applied, because the observations underlying the experience-based knowledge acquired are lacking. For example, to validate the greenhouse management rules formalised from expert knowledge, Tchamitchian et al. (2006) used a two-step method rather than a direct validation of the rules themselves, which was not possible. The first step involved checking that the application of these rules really did result in the desired pattern of behaviour in the greenhouse (as expressed when building the rules), without questioning the agronomic validity of this behaviour. The second step involved assessing the quality of production obtained by applying these rules, the goal being to obtain appropriate production levels from the greenhouse. Attempts at the direct validation of a given rule have only made explicit which pieces of agronomic knowledge can be used to support a given rule. However, it would not have been possible to design the rule from this identified scientific knowledge, generally because the scopes of the scientific knowledge and that of the lay expertise yielding the rule were different.

3. Methods for synthesizing information

The three main research methods currently used by agronomists (Fig. 1) are various types of field experiments, on-farm inquiries (e.g., Doré et al., 2008), and modelling (e.g., Rossing et al., 1997; Bergez et al., 2010). Field experiments provide validated knowledge meeting the scientific rules for data acquisition. This basic knowledge can be supplemented by inquiries providing data from real-world agricultural situations (farms). Modelling can be used to explore the response of key agronomic and environmental variables, such as, for example, yield or nitrogen loss, to climate, cropping system variables or societal changes. The data generated are then processed, mostly by classical methods, such as simulation studies, single-experiment data analysis, or group analysis. These methods could probably be complemented with two other methods: meta-analysis, involving the statistical synthesis of results from a series of studies, and comparative analyses of agroecosystems, involving the use of large-scale comparisons similar to those used in ecology (e.g., Fortunel et al., 2009).

3.1. Meta-analysis and agronomy

Meta-analysis (e.g., Borenstein et al., 2009) is more powerful than a simple narrative review of a series of studies, because it synthesises published data in a quantitative manner and makes it possible to assess the between-study variability of a variable of interest.

Both scientific researchers and decision-makers can benefit from meta-analysis in several ways (Sutton et al., 2000), as this approach provides a methodological framework for (i) exploring what has already been done on a given research topic and identifying more clearly where the gaps and uncertainties lie, (ii) generating an overview of divergent results, (iii) guiding decisions based on a systematic review and statistical analysis of all the available data related to a given topic, (iv) broadening the knowledge base and allowing replication for the testing of hypotheses, (v) adding to the cumulative development of science.

Most meta-analyses carried out to date have been performed in medical science (Normand, 1999; Sutton et al., 2000). This approach has been less systematically applied in other areas of research, such as ecology (e.g., Arnqvist and Wooster, 1995; Cardinale et al., 2006), and has sometimes been applied in agriculture (e.g., Bengtsson et al., 2005), animal science (Sauvant et al., 2008) and plant pathology (Rosenberg et al., 2004). In agronomy, meta-analysis methods have generally been used to compare the effects of different cropping techniques or of different cropping systems on yield or biomass production. For example, Miguez and Bollero (2005) used a meta-analysis method to summarise and describe quantitatively the effect of several winter cover crops on maize yield. The authors estimated the ratio of maize yield after a winter cover crop to maize yield with no cover from 37 published studies carried out in various regions of the USA and Canada. In another study, Miguez et al. (2008) studied the effects of planting density and nitrogen fertiliser on the biomass production of *Miscanthus × giganteus*, using 31 published studies including biomass measurements at different dates over several years. Drawing on published studies on sub-Saharan African agriculture, Chikowo et al. (2010) conducted a meta-analysis of factors controlling nitrogen and phosphorus capture and conversion efficiencies by major cereal crops. The meta-analysis carried out by Badgley et al. (2007) did not focus on a specific cropping technique, but was performed to compare two agricultural systems: organic versus conventional or low-intensity. The authors compared the yields obtained in an organic system with those obtained in conventional or low-intensity food production systems, based on yield data from 293 individual studies on various crops. These data were used to estimate the mean yield ratio for various food categories, for both developed and developing countries.

Diverse techniques for meta-analysis are available (e.g., Borenstein et al., 2009; Sutton et al., 2000), but meta-analysis should always include the following steps:

- i. Definition of the objective of the meta-analysis and of the variable of interest to be estimated from the data (e.g., in Miguez and Bollero, 2005, the variable of interest is the ratio of maize yield after a winter cover crop to maize yield in the absence of a cover crop).
- ii. Systematic review of the literature and/or of the dataset reporting values of the quantities of interest.
- iii. Analysis of data quality (i.e., quality of the experimental designs and of the measurement techniques).
- iv. Assessment of between-study variability and heterogeneity. Evaluation of the between-study variability of the variable of interest and of the heterogeneity of the accuracy of individual estimates is an important step in a meta-analysis and several statistical methods have been proposed to estimate between-

and within-study variances (Borenstein et al., 2009). Combination of the individual study estimates and estimation of a mean value for the variable of interest, for example, can be achieved by calculating a weighted sum of individual estimates derived from the studies collected in step ii.

- v. Assessment of publication bias. Publication bias occurs when only studies with highly significant results are published. In this case, a meta-analysis can lead to a biased conclusion and overestimation of the effect of a given factor. The 'funnel plot' technique can be used to deal with this issue (e.g., Borenstein et al., 2009).
- vi. Presentation of the results and of the level of uncertainty.

In the context of ecological intensification, the meta-analysis framework constitutes an interesting alternative to dynamic crop models. Dynamic crop models can be used both to assess the consequences of cropping techniques and environmental variables for crop production (e.g., Jones and Thornton, 2003) and to assess the effect of cropping systems on key environmental variables (e.g., Rolland et al., 2008), two key issues for ecological intensification. However, these models include several sources of uncertainty (Monod et al., 2006) and their predictions are not always reliable (e.g., Barbottin et al., 2008; Makowski et al., 2009). We believe that meta-analysis should be more systematically used by agronomists, to assess and compare the effects of cropping systems on productivity, risks of soil and water pollution, greenhouse gas emissions and biodiversity. A considerable body of experimental data is available for such purposes (e.g., Rochette and Janzen, 2005). Such data could be reviewed, combined and analysed with statistical techniques, to rank cropping systems as a function of their impact on key environmental variables, such as water nitrate content, greenhouse gas emissions (e.g., N₂O) and the presence/absence of species of ecological interest (e.g., earthworms, birds). However, meta-analysis requires the use of appropriate techniques and the value of a meta-analysis may be greatly decreased if the six steps outlined above are not rigorously implemented.

3.2. Comparative analysis of agroecosystems

Information useful for the ecological intensification of agroecosystems may be obtained from comparative analyses of the structural and functional properties and performance of contrasting agroecosystems. Similar approaches, based on temporal or spatial comparisons, are used in other fields of research, such as plant sciences (Wright et al., 2004; Vile et al., 2005; Mauseth, 2006), evolution sciences (Schluessel et al., 2008) and marine ecology (Fuhrman and Steele, 2008). The comparative analysis of agroecosystems and comparisons of agroecosystems with natural ecosystems involve the simultaneous analysis of multiple criteria, with evaluation of the extent to which they display specific system properties. Several approaches have been proposed for this purpose (e.g., Pannell and Glenn, 2000; de Bie, 2000; Xu and Mage, 2001; López-Ridaura et al., 2002; Giampietro, 2003), based largely on concepts formulated more than a decade ago, by authors such as Conway (1987) and Marten (1988). These methods evaluate indicators relating to the properties of agroecosystems, such as productivity, stability and resilience. These properties are often interdependent and, as pointed out by Marten (1988), they are not universal and must be redefined under each new set of conditions. As discussed above, studies of the local knowledge sustaining various mechanisms of indigenous resilience across contrasting agroecosystems, particularly at the scale of the landscape and its functionality (e.g., Birman et al., 2010), are also a promising starting point for obtaining information useful for ecological intensification. In the next few paragraphs, we examine briefly some critical issues relating to the choice of indicators in multicriteria

evaluations and identify innovative ways of looking at the relationship between structure and function in agroecosystems.

3.2.1. Comparative analysis based on multiple indicators

In practice, the implementation of multicriteria analytical frameworks often involves the selection of a number of indicators (or the use of a list of predetermined indicators) and of reference threshold values for each indicator. The selection of indicators is frequently biased towards the disciplinary standpoint of the observer or highly influenced by certain stakeholders, so 'quality control' methods for evaluating the choice of indicators are necessary. In their examination of the choice of indicators in different case studies, Groot and Pacini (2010) argued that multicriteria evaluations should involve the analysis of four main system properties: performance, diversity, coherence and connectedness, which can be approached from four dimensions: physical, ecological, productive and social. Performance relates to functional properties of the agroecosystem, such as capacity, stability and resilience. Diversity relates to the structural properties sustaining such functions. Indicators of coherence describe the degree of interaction between components or subsystems within an agroecosystem, and connectedness describes interactions with adjacent systems (i.e., other agroecosystems, urban or natural systems, etc.). When several indicators are considered simultaneously, it may be pertinent to check whether all the relevant criteria pertaining to system performance, diversity, coherence or connectedness are given equal importance. For example, López-Ridaura et al. (2002) and Pacini et al. (2003) used two sets of indicators in two independent evaluations of agroecosystems. Although both methods considered multiple criteria pertaining to system sustainability, they weighted the various system properties and/or dimensions of sustainability differently.

In general, comparative analyses based on indicators provide a static picture of the status of agroecosystems at one particular point in time, without considering the underlying feedback and system dynamics responsible for bringing the system to its current status and for any subsequent change to that status. Beyond comparing multiple indicators and the tradeoffs between them, the comparative analysis of agroecosystems should aim to distil the relationships between relevant properties; e.g., between performance on the one hand, and diversity, coherence and connectedness on the other. A common denominator of the indicators used in multi-criteria evaluations is their interdependence and their dependence on the structural diversity of the agroecosystem. This interdependence results from the co-adaptation of agroecosystem components over time. The structural diversity of agroecosystems, corresponding to the diversity of system components and their interrelationships, is only functional when organised in a specific way.

3.2.2. Analysing the structure and functioning of agroecosystems

It is often postulated that the ecological intensification of agroecosystems may be achieved through gradual diversification to capitalise on regulatory principles and mechanisms inherent to natural ecosystems (see above and, for example, Altieri, 1999; Gliessman, 2001; Wezel et al., 2009). Knowledge of the structural diversity of an agroecosystem, however, may not be sufficient to explain its behaviour, and the way in which the diverse components of the system relate to each other should also be known. Moreover, unnecessarily high degrees of diversity of system components and flows within systems with poorly organised configurations may lead to redundancy (Kauffman, 1995; Ulanowicz, 2004). Here, we examine some methods for studying the diversity and organisation of system components based on the theory of networks that may be used in the comparative analysis of agroecosystems.

Indicators of network complexity and organisation have been derived from communication science. They were first used in eco-

nomics by Leontief (1951, 1966), and later introduced into ecology by Hannon (1973). Indicators, such as average mutual information (AMI) and ascendency (A), were proposed by Ulanowicz (1997, 2004) for characterisation of the development capacity (in terms of increased organisation) of ecological systems, and have recently been used in comparative analyses of agroecosystems (Rufino et al., 2009). This approach is known as ecological network analysis, and Rufino et al. (2009) presented a set of indicators including AMI, A, and Finn's cycling index, for assessment of the diversity and organisation of system components governing N flows and food self-sufficiency in three smallholder crop-livestock systems from Ethiopia, Kenya and Zimbabwe. Farm systems are conceptualised as networks, with the household and the farming activities represented as compartments and the N flows represented as connections between compartments. In this example, indicators assessing network size, activity, cycling, organisation and diversity of the N flows were compared with indicators of productivity and household food self-sufficiency. This analysis revealed that although the amounts of N cycled were small and similar at all sites, resource use efficiency and dependence on external resources differed widely between these apparently 'comparable' agroecosystems. System performance was positively related to N flow network size, organisation and N cycling, consistent with the hypothesis that increasing the organisation of resource cycling within resource-limited agroecosystems may render these systems more adaptable and less vulnerable.

The main hypothesis underlying the use of these indicators is that agroecosystems retain the properties of the natural ecosystems for which these indices were derived. Ulanowicz (2004) calculated the value of several indicators of network size and organisation, such as the number of different nodes and flows, their roles and their connectivity, for a number of natural ecosystems and agroecosystems. This exercise revealed wider gaps between these systems in terms of indicators of organisation than for the magnitude of energy matter and information flow within them. In other words, increasing organisation makes it possible to do much more with the same resources, while contributing to system stability. The extent and the manner in which organisation contributes to building resilience in agroecosystems is a fascinating research area that remains largely unexplored. Existing frameworks of thinking about resilience in the field of ecology and nature conservation may also be of interest here (e.g., Walker et al., 2010). An indirect measurement of the organisation of an agroecosystem is its energy and entropy balance. Svirezhev (2000) proposed the use of thermodynamics concepts to assess the sustainability of agroecosystems, based on the principle that an ecosystem in equilibrium with its environment has a certain 'capacity' to absorb anthropogenic stress that is regulated by its capacity to expel entropy back towards the environment (the 'entropy pump'). This capacity, which emerges from various agroecosystem properties, can be used to characterise the status of an agroecosystem with respect to the adjacent natural ecosystem from which it has been derived.

Many of the properties of agroecosystems are often interdependent, together determining the vulnerability and adaptation capacity of these systems in the face of external shocks and stressors (Luers, 2005). Far from being postulates of a new theory, these properties are discussed here as operational, working concepts. We know that the provision of agroecosystem service functions is regulated by the intrinsic properties of these systems, the functionality of which can be influenced by design. In practical terms, 'design' implies proposing alternative configurations for the organisation of energy, matter and information flows towards, within and from the system in space and time. The examples examined here indicate that, up to a certain critical level, an increase in the diversity of system components and interrelationships confers desirable properties on agroecosystems consistent

with the paradigm of ecological intensification. However, these properties manifest themselves as patterns in space and time that become more evident at particular scales and are often described as variability and/or heterogeneity at other scales. Diversity and spatio-temporal variability or heterogeneity are inherent to agroecosystems (Burel and Baudry, 2003), and may represent constraints to the representation of these systems in prototyping or modelling, which is often based on modal agroecosystem configurations.

4. Overall discussion and conclusion

Wide new avenues seem to be opening up in agronomy to guide ecological intensification. We have tried here to identify new sources of knowledge and methods and to consider their potential role (Fig. 1). The analysis, use and optimisation of biological regulation in agroecosystems are the most commonly promoted methods of ecological intensification. This approach frequently involves enlarging the foundations of agronomic knowledge to cover biotic components of the system and their interactions. This ecological analysis of the whole system is of paramount importance, and further investment in this approach is required. This will involve the expansion of agronomic knowledge through classical avenues of research, involving the generation of data mostly through modelling and on-station experiments, and their analysis through simulation studies or statistical hypothesis testing. Our proposed approach is complementary to attempts to increase our understanding of biological regulations in agroecosystems and to use this knowledge for ecological intensification. Indeed, the extension of sources of knowledge to natural ecosystems and farmers' knowledge relates mostly to biological regulation and is fundamentally consistent with the scientific approach to acquiring knowledge about biological regulation in agroecosystems. The extension of sources of knowledge to the results of plant sciences research is more debatable. For example, Vanloqueren and Baret (2009) argued that genetic engineering closes off avenues of agroecological innovation. However, plant science results are not inevitably linked to a single technological regime. Agronomists, if they were aware of current knowledge in plant sciences, could make use of some of this knowledge to rebalance technological regimes or to construct new ones. The expansion of sources of knowledge will also indirectly promote ways of generating data that are little used at the moment. Most agronomic data are still acquired through on-station trials and modelling. The extension of sources of knowledge to farmers' knowledge and natural ecosystems will highlight alternative methods of data generation. This will, in turn, incite the development of new data processing methods, such as meta-analysis and comparative studies.

The new avenues outlined here will require major methodological investment. Indeed, the extension of sources of knowledge suggested here is far from straightforward. Plant science results must be thoroughly screened by groups of agronomists and plant scientists working together, to identify the most promising results for use in ecological intensification. Three major points should be made:

- (i) Most plant science knowledge of potential use in agronomy is based on genetic drivers. As gene expression depends on environmental conditions, the use of plant science data in ecological intensification will require qualification and quantification of the corresponding genotype \times environment interactions, for a range of cropping systems, soils and climatic conditions (see for example Spiertz et al., 2007).
- (ii) All dimensions of cropping system management may benefit from a greater knowledge of plant biology and soil ecology: crop rotation sequences, soil management, crop management,

etc. Furthermore, most of the issues raised by ecological intensification can be addressed: yield increase, cut-off for the use of limited resources through better mineral use efficiency, decrease in pesticide use through the adoption of new crop protection methods, etc.

- (iii) Our paper is limited to a few examples. To our knowledge, probably due to schism between agronomists and plant scientists, no formal attempt to enlarge this list has been made by systematically tracking plant science results of potential use in cropping system design. Such tracking of results and the publication of the findings obtained would nonetheless be of considerable interest.

The use of knowledge relating to natural ecosystems requires clarification concerning what to study and how, for each of the properties of agroecosystems that ecological intensification aims to improve. This suggests a possible step-wise course of action for agronomists seeking to mimic natural ecosystems:

- Selection of the functions agronomists wish to improve (for example, nutrient cycle management).
- Identification, in natural ecosystems, of the structural characteristics (spatial heterogeneity, diversification of vegetation strata, variability of species in time and space, etc.) modifying these functions.
- Definition of the qualitative or quantitative relationships linking properties and functions.
- Transposition of these functions to agricultural conditions.
- Use of these functions for the design of agroecosystems with specified aims.
- Checking that the new agroecosystems express the targeted functions and have no undesirable properties.

This procedure seems far more complex than simply trying to design agroecosystems "as similar as possible" to natural ecosystems.

Farmers' knowledge seems to be extremely valuable, and its use in association with scientific knowledge requires appropriate processing by methods that are not yet well established. Specific methods remain to be adapted from other domains or developed. The first methodological requirement is a more profound analysis of local knowledge to determine which processes (ecological or otherwise) should be selected and how they can be used or manipulated. Davis and Ruddle (2010) analysed the ways in which ecological knowledge (local, traditional or indigenous) is used and concluded that the same level of scrutiny as for scientific experimental results should be applied before such knowledge is accepted. However, this local knowledge is built within specific 'systems of knowledge' (Davis and Ruddle, 2010), and therefore cannot be analysed purely in terms of its content relevant to agronomy or ecological science. It must also be analysed from a social point of view (which processes lead to this knowledge? How is it shared, transmitted, etc.?). This analysis calls for pluridisciplinary approaches. We also need to design approaches inspired by or directly making use of the argumentation theory and methods developed in the domain of artificial intelligence (Amgoud and Prade, 2009).

The use of meta-analysis methods for ecological intensification benefits from extensive experience in other research areas, and follows guidelines that have proved to be effective. Nevertheless, data acquisition in agronomy has not traditionally been organised with the requirements of subsequent meta-analyses in mind. As a consequence, considerable effort is required to adapt the methods to existing agronomic data and to establish guidelines for the generation of further data. Finally, comparative studies in agriculture often remain descriptive, and are not always oriented to

identify the relationships between agroecosystem structure and functioning—undoubtedly a new challenge for agronomic research. Addressing this aim will require the development of guidelines for site selection, characterisation methods, data processing, etc.

Finally, each of the five topics outlined will probably require specific organisation within research institutes. They may also induce changes in academic curricula in agronomy, as plant scientists and agronomists currently follow different curricula, with little in the way of shared knowledge, concepts and technical skills.

Acknowledgement

We thank Alain Bône and Julie Sappa for their skilled assistance.

References

- Abbona, E.A., Sarandón, S.J., Marasas, M.E., Astier, M., 2007. Ecological sustainability evaluation of traditional management in different vineyard systems in Berisso, Argentina. *Agric. Ecosyst. Environ.* 119, 335–345.
- Acosta-Martinez, V., Dowd, S., Sun, Y., Allen, V., 2008. Tag-encoded pyrosequencing analysis of bacterial diversity in a single soil type as affected by management and land use. *Soil Biol. Biochem.* 40, 2762–2770.
- Aerts, R., 1999. Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks. *J. Exp. Bot.* 50, 29–37.
- Affholder, F., Jourdain, D., Morize, M., Quang, D.D., Ricome, A., 2008. Ecological intensification in the mountains of Vietnam: constraints to the adoption of cropping systems based on mulches and cover crops. *Cah. Agric.* 17, 289–296.
- Affholder, F., Jourdain, D., Quang, D.D., Tuong, T.P., Morize, M., Ricome, A., 2010. Constraints to farmers' adoption of direct-seeding mulch-based cropping systems: a farm scale modeling approach applied to the mountainous slopes of Vietnam. *Agric. Syst.* 103, 51–62.
- Alsinet, T., Cheshevar, C.I., Godo, L., Simari, G.R., 2008. A logic programming framework for possibilistic argumentation: formalization and logical properties. *Fuzzy Sets Syst.* 159, 1208–1228.
- Altieri, M.A., 2002. Agroecology: the science of natural resource management for poor farmers in marginal environments. *Agric. Ecosyst. Environ.* 93, 1–24.
- Altieri, M.A., 1999. The ecological role of biodiversity in agroecosystems. *Agric. Ecosyst. Environ.* 74, 19–31.
- Altieri, M.A., 1989. Agroecology—a new research and development paradigm for world agriculture. *Agric. Ecosyst. Environ.* 27, 37–46.
- Amgoud, L., Kaci, S., 2007. An argumentation framework for merging conflicting knowledge bases. *Int. J. Approx. Reason.* 45, 321–340.
- Amgoud, L., Prade, H., 2009. Using arguments for making and explaining decisions. *Artif. Intell.* 173, 413–436.
- Arnqvist, G., Wooster, D., 1995. Metaanalysis—synthesizing research findings in ecology and evolution. *Trends Ecol. Evol.* 10, 236–240.
- Baars, T., de Vries, A., 1999. Facilitating empirical learning in agriculture—scientific knowledge and practical choices: forging a link between the way of the researcher/extensionist and the way of the farmer. In: Zanoli, R., Krell, R. (Eds.), *Landscape Ecology in Agroecosystems Management Rome (ITA)*, vol. 58. REU Technical Series, pp. 49–56.
- Badgley, C., Moghtader, J., Quintero, E., Zakem, E., Chappell, M.J., Aviles-Vazquez, K., Samulon, A., Perfecto, I., 2007. Organic agriculture and the global food supply. *Renew. Agric. Food Syst.* 22, 86–108.
- Bais, H.P., Park, S.W., Weir, T.L., Callaway, R.M., Vivanco, J.M., 2004. How plants communicate using the underground information superhighway. *Trends Plant Sci.* 9, 26–32.
- Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S., Vivanco, J.M., 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* 57, 233–266.
- Ballard, H.L., Fernandez-Gimenez, M.E., Sturtevant, V.E., 2008. Integration of local ecological knowledge and conventional science: a study of seven community-based forestry organizations in the USA. *Ecol. Soc.* 13 (art. n. 37), <http://www.ecologyandsociety.org/vol13/iss2/art37/>.
- Barbottin, A., Makowski, D., Le Bail, M., Jeuffroy, M.-H., Bouchard, C., Barrier, C., 2008. Comparison of models and indicators for categorizing soft wheat fields according to their grain protein contents. *Eur. J. Agron.* 29, 175–183.
- Barzman, M.S., Mills, N.J., Thi Thu Cuc, N., 1996. Traditional knowledge and rationale for weaver ant husbandry in the Mekong delta of Vietnam. *Agric. Human Values* 13, 2–9.
- Bench-Capon, T.J.M., Dunne, P.E., 2007. Argumentation in artificial intelligence. *Artif. Intell.* 171, 619–641.
- Bengtsson, J., Ahnström, J., Weibull, A.-C., 2005. The effects of organic agriculture on biodiversity and abundance: a meta-analysis. *J. Appl. Ecol.* 42, 261–269.
- Bergez, J.-E., Colbach, N., Crespo, O., Garcia, F., Jeuffroy, M.-H., Justes, E., Loyce, C., Munier-Jolain, N., Sadok, W., 2010. Designing crop management systems by simulation. *Eur. J. Agron.* 32, 3–9.
- Berkes, F., 1999. *Sacred Ecology: Traditional Ecological Knowledge and Resource Management*. Taylor and Francis, Philadelphia, VA (USA).
- Bertin, C., Yang, X.H., Weston, L.A., 2003. The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil* 256, 67–83.
- Birman, D., Moraine, M., Tiftonell, P., Martin, P., Clouvel, P., 2010. Ecosystem services assessment in complex agricultural landscapes using farmers' perception. In: *Proceedings of AGRO2010, The Scientific International Week on Agronomy*, Montpellier (FRA).
- Bommel, P., Bonaudo, T., Barbosa, T., Bastos da Veiga, J., Vieira Pak, M., Tourrand, J.-F., 2010. The complex relationship between cattle ranching and the forest in Brazilian Amazonia. A multiagent modelling approach. *Cah. Agric.* 19, 104–111.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T., Rothstein, H.R., 2009. *Introduction to Meta-Analysis*. John Wiley and Sons, Chichester (GBR).
- Boss, P.K., Bastow, R.M., Mylne, J.S., Dean, C., 2004. Multiple pathways in the decision to flower: enabling, promoting, and resetting. *Plant Cell* 16, S18–S31.
- Bowen, G.D., Kovira, A.D., 1999. The rhizosphere and its management to improve plant growth. *Adv. Agron.* 66, 1–102.
- Brookfield, H., Parsons, H., Brookfield, M., 2003. *Agrodiversity: Learning from Farmers Across the World*. United Nations University Press, New-York, NY (USA).
- Brussaard, L., de Ruiter, P.C., Brown, G.G., 2007. Soil biodiversity for agricultural sustainability. *Agric. Ecosyst. Environ.* 121 (3), 233–244.
- de Bruxelles, G.L., Roberts, M.R., 2001. Signals regulating multiple responses to wounding and herbivores. *Crit. Rev. Plant Sci.* 20, 487–521.
- Burel, F., Baudry, J., 2003. *Landscape Ecology: Concepts, Methods and Applications*. Science Publishers, Inc., Enfield, NH, USA.
- Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M., Jouseau, C., 2006. Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature* 443, 989–992.
- Cassman, K.G., 1999. Ecological intensification of cereal production systems: yield potential, soil quality, and precision agriculture. *Proc. Natl. Acad. Sci. U.S.A.* 96, 5952–5959.
- Chalmers, N., Fabricius, C., 2007. Expert and generalist local knowledge about land-cover change on South Africa's wild coast: can local ecological knowledge add value to science? *Ecol. Soc.* 12 (art. n. 10), <http://www.ecologyandsociety.org/vol12/iss1/art10/>.
- Chevassus au Louis, B., Griffon, M., 2008. La nouvelle modernité: une agriculture productive à haute valeur écologique. *Déméter: Économie et Stratégies Agricoles* 14, 7–48.
- Chikowo, R., Corbeels, M., Mapfumo, P., Tiftonell, P., Vanlauwe, B., Giller, K.E., 2010. Benefits of integrated soil fertility and water management in semi-arid West Africa: an example study in Burkina Faso. *Nutrient Cycl. AgroEcosyst.* 88, 59–77.
- Conway, G.R., 1987. The properties of agroecosystems. *Agric. Syst.* 24, 95–117.
- Cornelissen, A.M.G., van den Berg, J., Koops, W.J., Kaymak, U., 2003. Elicitation of expert knowledge for fuzzy evaluation of agricultural production systems. *Agric. Ecosyst. Environ.* 95, 1–18.
- Davis, A., Ruddle, K., 2010. Constructing confidence: rational skepticism and systematic enquiry in local ecological knowledge research. *Ecol. Appl.* 20, 880–894.
- Dawson, T., Fry, R., 1998. Agriculture in nature's image. *Trends Ecol. Evol.* 13, 50–51.
- Debaeke, P., Munier-Jolain, N., Bertrand, M., Guichard, L., Nolot, J.-M., Faloya, V., Saulas, P., 2009. Iterative design and evaluation of rule-based cropping systems: methodology and case studies. A review. *Agron. Sustain. Dev.* 29, 73–86.
- de Bie, C.A.J.M., 2000. Comparative performance analysis of agro-ecosystems. Ph.D. thesis. Wageningen University, Wageningen (NLD).
- dePury, D.G.G., Farquhar, G.D., 1997. Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant Cell Environ.* 20, 537–557.
- de Soto, H., 2000. *The Mystery of Capital: Why Capitalism Triumphs in the West and Fails Everywhere Else*. Basic Books, New-York, NY (USA).
- Doré, T., Clermont-Dauphin, C., Crozat, Y., David, C., Jeuffroy, M.H., Loyce, C., Makowski, D., Malézieux, E., Meynard, J.M., Valantin-Morison, M., 2008. Methodological progress in on-farm regional agronomic diagnosis. A review. *Agron. Sustain. Dev.* 28, 151–161.
- Dudareva, N., Pichersky, E., 2008. Metabolic engineering of plant volatiles. *Curr. Opin. Biotechnol.* 19, 181–189.
- Egger, K., 1986. Ecological intensification. Soil conservation and improvement of tropical soils by pastoral agroforestry systems. *Collection Documents Systèmes Agraires* 6, 129–135.
- Ewel, J., 1999. Natural systems as models for the design of sustainable systems of land use. *Agric. Syst.* 45, 1–21.
- FAO, 2009. *Glossary on Organic Agriculture*. FAO, Rome (ITA).
- Fortunel, C., Garnier, E., Joffre, R., Kazakou, E., Qested, H., Grigulis, K., Lavorel, S., Ansquer, P., Castro, H., Cruz, P., Dolezal, J., Eriksson, O., Freitas, H., Golodets, C., Jouany, C., Kigel, J., Kleyer, M., Lehsten, V., Leps, J., Meier, T., Pakeman, R., Papadimitriou, M., Papanastasis, V.P., Quetier, F., Robson, M., Sternberg, M., Theau, J.-P., Thebault, A., Zarovali, M., 2009. Leaf traits capture the effects of land use changes and climate on litter decomposability of grasslands across Europe. *Ecology* 90, 598–611.
- Friedman, D.B., Kanwat, C.P., Headrick, M.L., Patterson, N.J., Neely, J.C., Smith, L.U., 2007. Importance of prudent antibiotic use on dairy farms in South Carolina: a pilot project on farmers' knowledge, attitudes and practices. *Zoonoses Public Health* 54, 366–375.
- Fuhrman, J.A., Steele, J.A., 2008. Community structure of marine bacterioplankton: patterns, networks, and relationships to function. *Aquat. Microb. Ecol.* 53, 69–81.
- Fukai, S., 1993. Intercropping - bases of productivity - introduction. *Field Crops Res.* 34, 239–245.
- Funes-Monzote, F.R., Monzote, M., Lantinga, E.A., van Keulen, H., 2009. Conversion of specialised dairy farming systems into sustainable mixed farming systems in Cuba. *Environ. Dev. Sustain.* 11, 765–783.

- Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R., Martinelli, L.A., Seitzinger, S.P., Sutton, M.A., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320, 889–892.
- Giampietro, M., 2003. Complexity and scales: the challenge for integrated assessment. In: Rotmans, J., Rothman, D. (Eds.), *Scaling in Integrated Assessment*. Swets & Zeitlinger, Exton, PA, pp. 293–327.
- Glass, A.D.M., 2003. Nitrogen use efficiency of crop plants: physiological constraints upon nitrogen absorption. *Crit. Rev. Plant Sci.* 22, 453–470.
- Gliessman, S.R., 2001. *Agroecology: Ecological Processes in Sustainable Agriculture*. Lewis Publisher, Boca Raton, FL (USA).
- Grace, D., Randolph, T., Affognon, H., Dramane, D., Diall, O., Clausen, P.-H., 2009. Characterisation and validation of farmers' knowledge and practice of cattle trypanosomiasis management in the cotton zone of West Africa. *Acta Trop.* 111, 137–143.
- Groot, J.C.J., Pacini, G.C., 2010. Evaluation of indicators sets using a polyocular perspective on agroecosystems. In: Dogliotti, S., Rossing, W.A.H., Cittadini, E., Albín, A. (Eds.), *Proceedings of the International Congress on Co-innovation in Sustainable Livelihood Systems*. Lavalleja, Uruguay.
- Grossman, A., Takahashi, H., 2001. Macronutrient utilization by photosynthetic eukaryotes and the fabric of interactions. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52, 163–210.
- Grossman, J.M., 2003. Exploring farmer knowledge of soil processes in organic coffee systems of Chiapas, Mexico. *Geoderma* 111, 267–287.
- Gunderson, L.H., Holling, C.S. (Eds.), 2002. *Panarchy: Understanding Transformations in Human and Natural Systems*. Island Press, Washington, DC.
- Hannon, B., 1973. The structure of ecosystems. *J. Theor. Biol.* 41, 535–546.
- Hatfield, C.B., 1997. Oil back on the global agenda. *Nature* 387, 121.
- He, X.H., Critchley, C., Bledsoe, C., 2003. Nitrogen transfer within and between plants through common mycorrhizal networks (CMNs). *Crit. Rev. Plant Sci.* 22, 531–567.
- Henderson, J., 2010. Expert and lay knowledge: a sociological perspective. *Nutr. Dietet.* 67, 4–5.
- Holling, C.S., 1973. Resilience and stability of ecological systems. *Annu. Rev. Ecol. Syst.* 4, 1–23.
- Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J., Wardle, D.A., 2005. Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* 75, 3–35.
- Hubert, B., Rosegrant, M., van Boekel, M.A.J.S., Ortiz, R., 2010. The future of food: scenarios for 2050. *Crop Sci.* 50, S33–S50.
- Hughes, G., Madden, L.V., 2002. Some methods for eliciting expert knowledge of plant disease epidemics and their application in cluster sampling for disease. *Crop Prot.* 21, 203–215.
- Jackson, L.E., Burger, M., Cavanaugh, T.R., 2008. Roots, nitrogen transformations, and ecosystem services. *Annu. Rev. Plant Biol.* 59, 341–363.
- Jackson, W., 2002. Natural systems agriculture: a truly radical alternative. *Agric. Ecosyst. Environ.* 88, 111–117.
- Jackson, W., Jackson, L.L., 1999. Developing high seed yielding perennial polycultures as a mimic of mid-grass prairie. In: Lefroy, E.C., Hobbs, R.J., O'Connor, M.H., Pate, J.S. (Eds.), *Agriculture as a Mimic of Natural Ecosystems (Current Plant Science and Biotechnology in Agriculture, vol. 37)*. Kluwer Academic Publishing, Dordrecht (NLD), pp. 1–38.
- Jones, P.G., Thornton, P.K., 2003. The potential impacts of climate change on maize production in Africa and Latin-America in 2055. *Glob. Environ. Change* 13, 51–59.
- Kauffman, S., 1995. *At Home in the Universe: The Search for Laws of Self-Organization and Complexity*. Oxford University Press, 321 pp.
- Kessler, A., Baldwin, I.T., 2002. Plant responses to insect herbivory: the emerging molecular analysis. *Annu. Rev. Plant Biol.* 53, 299–328.
- Kiers, E.T., Denison, R.F., 2008. Sanctions, cooperation, and the stability of plant-rhizosphere mutualisms. *Annu. Rev. Ecol. Syst.* 39, 215–236.
- Lançon, J., Wery, J., Rapidel, B., Angokaye, M., Gerardeaux, E., Gaborel, C., Ballo, D., Fadegnon, B., 2007. An improved methodology for integrated crop management systems. *Agron. Sustain. Dev.* 27, 101–110.
- Le Bot, J., Bénard, C., Robin, C., Bourgaud, F., Adamowicz, S., 2009. The 'trade-off' between synthesis of primary and secondary compounds in young tomato leaves is altered by nitrate nutrition: experimental evidence and model consistency. *J. Exp. Bot.* 60, 4301–4314.
- Lefroy, E.C., Hobbs, R.J., O'Connor, M.H., Pate, J.S., 1999. *Agriculture as a Mimic of Natural Ecosystems*. Kluwer Academic Publishing, Dordrecht (NLD).
- Lemanceau, P., Expert, D., Gaymard, F., Bakker, P.A.H.M., Briat, J.-F., 2009. Role of iron in plant-microbe interactions. In: *Plant Innate Immunity, Advances in Botanical Research*, vol. 51, pp. 491–549.
- Leontief, W.W., 1966. *Input-Output Economics*. Oxford University Press, New York, USA.
- Leontief, W.W., 1951. *The Structure of the American Economy, 1919–1939: An Empirical Application of Equilibrium Analysis*. Oxford University Press.
- Ley, T., Kump, B., Albert, D., 2010. A methodology for eliciting, modelling, and evaluating expert knowledge for an adaptive work-integrated learning system. *Int. J. Human-Computer Studies* 68, 185–208.
- Li, H., Shen, J., Zhang, F., Clairrotte, M., Drevon, J.J., Le Cadre, E., Hinsinger, P., 2008. Dynamics of phosphorus fractions in the rhizosphere of common bean (*Phaseolus vulgaris* L.) and durum wheat (*Triticum turgidum* durum L.) grown in monocropping and intercropping systems. *Plant Soil* 312, 139–150.
- Long, A.J., Nair, P.K.R., 1999. Trees outside forests: agro-, community, and urban forestry. *New Forests* 17 (1–3), 145–174.
- Lough, T.J., Lucas, W.J., 2006. Integrative plant biology: role of phloem long-distance macromolecular trafficking. *Annu. Rev. Plant Biol.* 57, 203–232.
- López-Ridaura, S., Masera, O., Astier, M., 2002. Evaluating the sustainability of complex socio-environmental systems. The MESMIS framework. *Ecol. Indic.* 2, 135–148.
- Luers, A.L., 2005. The surface of vulnerability: an analytical framework for examining environmental change. *Glob. Environ. Change-Human Policy Dimens.* 15, 214–223.
- Main, A.R., 1999. How much biodiversity is enough? *Agrofor. Syst.* 45, 23–41.
- Makowski, D., Tichit, M., Guichard, L., van Keulen, H., Beaudoin, N., 2009. Measuring the accuracy of agro-environmental indicators. *J. Environ. Manage.* 90, S139–S146.
- Malézieux, E., 2011. Designing cropping systems from Nature. *Agron. Sustain. Dev.*, in press, doi:10.1007/s13593-011-0027-z.
- Malézieux, E., Crozat, Y., Dupraz, C., Laurans, M., Makowski, D., Ozier-Lafontaine, H., Rapidel, B., de Tourdonnet, S., Valantin-Morison, M., 2009. Mixing plant species in cropping systems: concepts, tools and models. A review. *Agron. Sustain. Dev.* 29, 43–62.
- Marten, G.G., 1988. Productivity, Stability, Sustainability, Equitability and Autonomy as Properties for Agroecosystem Assessment. *Agric. Sys.* 26, 291–316.
- Matson, P.A., Parton, W.J., Power, A.G., Swift, M.J., 1997. Agricultural intensification and ecosystem properties. *Science* 277, 504–509.
- Maurel, C., Verdoucq, L., Luu, D.-T., Santoni, V., 2008. Plant aquaporins: membrane channels with multiple integrated functions. *Annu. Rev. Plant Biol.* 59, 595–624.
- Mauseth, J.D., 2006. Structure-function relationships in highly modified shoots of Cactaceae. *Ann. Bot.* 98, 901–926.
- McNeely, J.A., Scherr, S.J., 2003. *Ecogriculture, Strategies to Feed the World and Save Wild Biodiversity*. Island Press, Washington, DC (USA).
- McSteen, P., Leysner, O., 2005. Shoot branching. *Annu. Rev. Plant Biol.* 56, 353–374.
- Médiéne, S., Valantin-Morison, M., Sarthou, J.-P., de Tourdonnet, S., Gosme, M., Bertrand, M., Roger-Estrade, J., Aubertot, J.-N., Rusch, A., Motisi, N., Pelosi, C., Doré, T., 2011. Agroecosystem management and biotic interactions. A review. *Agron. Sustain. Dev.*, in press, doi:10.1007/s13593-011-0009-1.
- Mehboob, I., Naveed, M., Zahir, Z.A., 2009. Rhizobial association with non-legumes: mechanisms and applications. *Crit. Rev. Plant Sci.* 28, 432–456.
- Meinen-Dick, R., DiGregorio, M., McCarthy, N., 2004. Methods for studying collective action in rural development. *Agric. Syst.* 82, 197–214.
- Micallef, S.A., Shiaris, M.P., Colon-Carmona, A., 2009. Influence of *Arabidopsis thaliana* accessions on rhizobacterial communities and natural variation in root exudates. *J. Exp. Bot.* 60, 1729–1742.
- Michon, G., de Foresta, H., Levang, P., Verdeaux, F., 2007. Domestic forests: a new paradigm for integrating local communities' forestry into tropical forest science. *Ecol. Soc.* 12 (art. n. 1), http://www.ecologyandsociety.org/vol12/iss2/art1/.
- Michon, G., de Foresta, H., Levang, P., 1995. Strategies agroforestières et développement durable: les agroforêts à Damar de Sumatra. *Nat. Sci. Soc.* 3, 207–221.
- Miguez, F.E., Bollero, G.A., 2005. Review of corn yield response under winter cover cropping systems using meta-analytic methods. *Crop Sci.* 45, 2318–2329.
- Miguez, F.E., Villamil, M.B., Long, S.P., Bollero, G.A., 2008. Meta-analysis of the effects of management factors on *Miscanthus × giganteus* growth and biomass production. *Agric. For. Meteorol.* 148, 1280–1292.
- Mikolasek, O., Khuyen, T.D., Medoc, J.-M., Porphyre, V., 2009. The ecological intensification of an integrated fish farming model: Recycling pig effluents from farms in Thai Binh province (North Vietnam). *Cah. Agric.* 18, 235–241.
- Monod, H., Naud, C., Makowski, D., 2006. Uncertainty and sensitivity analysis for crop models. In: Wallach, D., Makowski, D., Jones, J.W. (Eds.), *Working with Dynamic Crop Models*. Elsevier Edition, pp. 55–100.
- Monteith, J.L., 1977. Climate and efficiency of crop production in Britain. *Philos. Trans. R. Soc. London B* 281, 277–284.
- Nair, P.K.R., 2001. Do tropical homegardens elude science, or is it the other way around? *Agrofor. Syst.* 53, 239–245.
- Navarrete, M., Jeannequin, B., Sebillotte, M., 1997. Vigour of greenhouse tomato plants (*Lycopersicon esculentum* Mill.): analysis of the criteria used by growers and search for objective criteria. *J. Hortic. Sci.* 72, 821–829.
- van Noordwijk, M., Ong, C.K., 1999. Can the ecosystem mimic hypotheses be applied to farms in African savannahs? *Agrofor. Syst.* 45, 131–158.
- Normand, S.-L.T., 1999. Tutorial in biostatistics. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat. Med.* 18, 321–359.
- Okoba, B.O., de Graaff, J., 2005. Farmers' knowledge and perceptions of soil erosion and conservation measures in the Central Highlands, Kenya. *Land Degrad. Devel.* 16, 475–487.
- Pacini, C., Wossink, A., Giesen, G., Vazzana, C., Huirne, R., 2003. Evaluation of sustainability of organic, integrated and conventional farming systems: a farm and field-scale analysis. *Agric. Ecosyst. Environ.* 95, 273–288.
- Pannell, D.J., Glenn, N.A., 2000. A framework for the economic evaluation and selection of sustainability indicators in agriculture. *Ecol. Econ.* 33, 135–149.
- Pietikäinen, A., Kytöviita, M.-M., 2007. Defoliation changes mycorrhizal benefit and competitive interactions between seedlings and adult plants. *J. Ecol.* 95, 639–647.
- Prior, L., 2003. Belief, knowledge and expertise: the emergence of the lay expert in medical sociology. *Sociol. Health. Ill.* 25, 41–57.
- Reed, M.S., Dougill, A.J., Taylor, M.J., 2007. Integrating local and scientific knowledge for adaptation to land degradation: Kalahari rangeland management options. *Land Degrad. Dev.* 18, 249–268.
- Reis, V.M., Baldani, J.I., Baldani, V.L.D., Dobereiner, J., 2000. Biological dinitrogen fixation in gramineae and palm trees. *Crit. Rev. Plant Sci.* 19, 227–247.

- Roberts, J.A., Elliott, K.A., Gonzalez-Carranza, Z.H., 2002. Abscission, dehiscence, and other cell separation processes. *Annu. Rev. Plant Biol.* 53, 131–158.
- Rochette, P., Janzen, H.H., 2005. Towards a revised coefficient for estimating N₂O emissions from legumes. *Nutrient Cycl. AgroEcosyst.* 73, 171–179.
- Rolland, M.-N., Gabrielle, B., Laville, P., Serça, D., Cortinovis, J., Larmanou, E., Lehuger, S., Cellier, P., 2008. Modeling of nitric oxide emissions from temperate agricultural soils. *Nutrient Cycl. AgroEcosyst.* 80, 75–93.
- Rosenberg, M.S., Garrett, K.A., Su, Z., Bowden, R.L., 2004. Meta-analysis in plant pathology: synthesizing research results. *Phytopathology* 94, 1013–1017.
- Rossing, W.A.H., Meynard, J.-M., van Ittersum, M.K., 1997. Model-based explorations to support development of sustainable farming systems: case studies from France and the Netherlands. *Eur. J. Agron.* 7, 271–283.
- Rufino, M.C., Tittonell, P., Reidsma, P., López-Ridaura, S., Hengsdijk, H., Giller, K.E., Verhagen, A., 2009. Network analysis of N flows and food self-sufficiency—a comparative study of crop-livestock systems of the highlands of East and southern Africa. *Nutrient Cycl. AgroEcosyst.* 85, 169–186.
- Ryan, P.R., Delhaize, E., Jones, D.L., 2001. Function and mechanism of organic anion exudation from plant roots. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52, 527–560.
- Ryan, P.R., Dessaux, Y., Thomashow, L.S., Weller, D.M., 2009. Rhizosphere engineering and management for sustainable agriculture. *Plant Soil* 321, 363–383.
- Saito, K., Linquist, B., Keobualapha, B., Shiraiwa, T., Horie, T., 2006. Farmers' knowledge of soils in relation to cropping practices: a case study of farmers in upland rice based slash-and-burn systems of northern Laos. *Geoderma* 136, 64–74.
- Sanchez, P.A., 1995. Science in agroforestry. *Agrofor. Syst.* 30, 5–55.
- Sanon, A., Andrianjaka, Z.N., Prin, Y., Bally, R., Thioulouse, J., Comte, G., Duponnois, R., 2009. Rhizosphere microbiota interferes with plant-plant interactions. *Plant Soil* 321, 259–278.
- Sauvant, D., Schmidely, P., Daudin, J.J., St-Pierre, N.R., 2008. Meta-analyses of experimental data in animal nutrition. *Animal* 2, 1203–1214.
- Schlüssel, V., Bennett, M.B., Bleckmann, H., Blomberg, S., Collin, S.R., 2008. Morphometric and ultrastructural comparison of the olfactory system in elasmobranchs: the significance of structure-function relationships based on phylogeny and ecology. *J. Morphol.* 269, 1365–1386.
- Schroth, G., da Fonseca, G.A.B., Harvey, C.A., Gascon, C., Vasconcelos, H.L., Izac, A.M.N., 2004. *Agroforestry and Biodiversity Conservation in Tropical Landscapes*. Island Press, Washington, DC (USA).
- Schroth, G., Lehmann, J., Rodrigues, M.R.L., Barros, E., Macedo, J.L.V., 2001. Plant-soil interactions in multistrata agroforestry in the humid tropics. *Agrofor. Syst.* 53, 85–102.
- Schulte, R.P.O., Lantinga, E.A., Struik, P.C., 2003. Analysis of the production stability of mixed grasslands I: a conceptual framework for the qualification of production stability in grassland ecosystems. *Ecol. Model.* 159, 43–69.
- Singh, R.K., Sureja, A.K., 2008. Indigenous knowledge and sustainable agricultural resources management under rainfed agro-ecosystem. *Indian J. Tradit. Knowl.* 7, 642–654.
- Sinzogan, A.A.C., van Huis, A., Kossou, D.K., Jiggins, J., Vodouhe, S., 2004. Farmers' knowledge and perception of cotton pests and pest control practices in Benin: results of a diagnostic study. *NJAS-Wagen J. Life Sci.* 52, 285–303.
- Spiertz, J.H.J., 2010. Nitrogen, sustainable agriculture and food security. A review. *Agron. Sustain. Dev.* 30, 43–55.
- Spiertz, J.H.J., Struik, P.C., van Laar, H.H., 2007. *Scale and Complexity in Plant Systems Research: Gene-Plant-Crop Relations*. Springer, 332 pp.
- Spitters, C.J.T., 1986. Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis. 1. Calculation of canopy photosynthesis. *Agric. Forest Meteorol.* 38 (1–3), 231–242.
- Spitters, C.J.T., Toussaint, H.A.J.M., Goudriaan, J., 1986. Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis. 1. Components of incoming radiation. *Agric. Forest Meteorol.* 38 (1–3), 217–229.
- Sprent, J.I., Sutherland, J.M., Defaria, S.M., 1987. Some aspects of the biology of nitrogen-fixing organisms. *Philos. Trans. R. Soc. Lond. Ser. B-Biol. Sci.* 317, 111–129.
- Stacey, G., Libault, M., Brechenmacher, L., Wan, J.R., May, G.D., 2006. Genetics and functional genomics of legume nodulation. *Curr. Opin. Plant Biol.* 9, 110–121.
- Steiner, K.G., 1998. Using farmers' knowledge of soils in making research results more relevant to field practice: experiences from Rwanda. *Agric. Ecosyst. Environ.* 69, 191–200.
- Sturz, A.V., Nowak, J., 2000. Endophytic communities of rhizobacteria and the strategies required to create yield enhancing associations with crops. *Appl. Soil Ecol.* 15, 183–190.
- Sutton, A.J., Abrams, K.R., Jones, D.R., Sheldon, T.A., Song, F., 2000. *Methods for Meta-analysis in Medical Research*. John Wiley, London (BGR).
- Swirezhev, Y.M., 2000. Thermodynamics and ecology. *Ecol. Model.* 132, 11–22.
- Swinton, S., Lupi, F., Robertson, G.P., Hamilton, S.K., 2007. Ecosystem services and agriculture: Cultivating agricultural ecosystems for diverse benefits. *Ecol. Econ.* 64, 245, 252.
- Tardieu, F., Tuberosa, R., 2010. Dissection and modeling of abiotic stress tolerance in plants. *Curr. Opin. Plant Biol.* 13 (2), 206–212.
- Tchamitchian, M., Martin-Clouaire, R., Lagier, J., Jeannequin, B., Mercier, S., 2006. Ser-riste: a daily set point determination software for glasshouse tomato production. *Comput. Electron. Agric.* 50, 25–47.
- Tittonell, P., 2007. Msimu wa Kupanda: targeting resources within diverse, heterogeneous and dynamic farming systems of East Africa. Ph.D. thesis. Wageningen University, Wageningen (NLD).
- Tittonell, P., van Wijk, M.T., Herrero, M., Rufino, M.C., de Ridder, N., Giller, K.E., 2009. Beyond resource constraints—exploring the biophysical feasibility of options for the intensification of smallholder crop-livestock systems in Vihiga district, Kenya. *Agric. Syst.* 101, 1–19.
- Torquebiau, E., 2007. *L'agroforesterie: des arbres et des champs*. L'Harmattan, Paris (FRA).
- Trenbath, B.R., 1993. Intercropping for the management of pests and diseases. *Field Crops Res.* 34, 381–405.
- Ulanowicz, R.E., 2004. Quantitative methods for ecological network analysis. *Comput. Biol. Chem.* 28, 321–339.
- Ulanowicz, R.E., 1997. *Ecology, the Ascendent Perspective*. Columbia University Press, New-York, NY (USA).
- Vandemeer, J.H., 2003. *Tropical Agroecosystems*. CRC Press, Boca Raton, FL (USA).
- Vanloqueren, G., Baret, P., 2009. How agricultural research systems shape a technological regime that develops genetic engineering but locks out agroecological innovations. *Res. Policy* 38 (6), 971–983.
- van der Heijden, M.G.A., Horton, T.R., 2009. Socialism in soil? The importance of mycorrhizal fungal networks for facilitation in natural ecosystems. *J. Ecol.* 97, 1139–1150.
- Vereijken, P., 1997. A methodical way of prototyping integrated and ecological arable farming systems (I/EAFS) in interaction with pilot farms. *Eur. J. Agron.* 7, 235–250.
- Vile, D., Garnier, E., Shipley, B., Laurent, G., Navas, M.L., Roumet, C., Lavorel, S., Diaz, S., Hodgson, J.G., Lloret, F., Midgley, G.F., Poorter, H., Rutherford, M.C., Wilson, P.J., Wright, I.J., 2005. Specific leaf area and dry matter content estimate thickness in laminar leaves. *Ann. Bot.* 96, 1129–1136.
- Walker, B., Sayer, J., Andrew, N.L., Campbell, B., 2010. Should enhanced resilience be an objective of natural resource management research for developing countries? *Crop Sci.* 50 (2), S10–S19.
- Walter, A., Silk, W.K., Schurr, U., 2009. Environmental Effects on Spatial and Temporal Patterns of Leaf and Root Growth. *Ann. Rev. Plant Biol.* 60, 279–304.
- Wang, Y., Li, J., 2008. Molecular basis of plant architecture. *Annu. Rev. Plant Biol.* 59, 253–279.
- Welbaum, G.E., Sturz, A.V., Dong, Z.M., Nowak, J., 2004. Managing soil microorganisms to improve productivity of agro-ecosystems. *Crit. Rev. Plant Sci.* 23, 175–193.
- Wezel, A., Bellon, S., Doré, T., Francis, C., Vallod, D., David, C., 2009. Agroecology as a science, a movement and a practice. A review. *Agron. Sustain. Dev.* 29, 503–515.
- Wissuwa, M., Mazzola, M., Picard, C., 2009. Novel approaches in plant breeding for rhizosphere-related traits. *Plant Soil* 321, 409–430.
- Witt, C., Fairhurst, T.H., Griffiths, W., 2006. The need to increase profitability in oil palm plantations: matching crop and nutrient management principles with evolving strategies. *Planter* 82, 377–403.
- Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias, J., Hikosaka, K., Lamont, B.B., Lee, T., Lee, W., Lusk, C., Midgley, J.J., Navas, M.L., Niinemets, U., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I., Roumet, C., Thomas, S.C., Tjoelker, M.G., Veneklaas, E.A.V.R., 2004. The worldwide leaf economics spectrum. *Nature* 428, 821–827.
- Xu, W., Mage, J.A., 2001. A review of concepts and criteria for assessing agroecosystem health including a preliminary case study of southern Ontario. *Agric. Ecosyst. Environ.* 83, 215–233.
- Zhang, W., Ricketts, T.H., Kremen, C., Carney, K., Swinton, S.M., 2007. Ecosystem services and dis-services to agriculture. *Ecol. Econ.* 64, 253–260.
- Zhang, L., van der Werf, W., Bastiaans, L., Zhang, S., Li, B., Spiertz, J.H.J., 2008. Light interception and utilization in relay intercrops of wheat and cotton. *Field Crops Res.* 107, 29–42.
- Zhu, X.-G., Long, S.P., Ort, D.R., 2010. Improving photosynthetic efficiency for greater yield. *Annual Rev. Plant Biol.* 61, 235–261.